arXiv:1508.00999v1 [math.NA] 5 Aug 2015

Approximation properties by generalized-Baskakov-Kantorovich-Stancu type operators

Abdul wafi, Nadeem Rao*

Department of Mathematics, Jamia Millia Islamia, New Delhi-110025, India

---

Abstract

In this paper, we introduce generalized Baskakov Kantorovich Stancu type operators and investigate direct result, local approximation and weighted approximation properties of these operators. Modulus of continuity, second modulus of continuity, Peeter's K-functional, weighted modulus of continuity and Lipschitz class are considered to prove our results.

Keywords: Baskakov Kantorovich operators, local approximation, second modulus of continuity, Peetre's K-functional, weighted modulus of continuity, Lipschitz space.
2010 Mathematics Subject Classification: 41A36, 41A25.

---

1. Introduction

For $f \in C[0,1]$, Bernstein[3] defined the linear positive operators which are the classical example of linear approximation as

$$B_n(f;x) = \sum_{k=0}^{n} \binom{n}{k} P_{n,k}(x) f\left(\frac{k}{n}\right), \tag{1}$$

where $P_{n,k}(x) = x^k(1-x)^{n-k}, x \in [0,1]$ and $n \in N$. In this paper, Bernstein showed that these operators approximate uniformly on [0,1] to every continuous function $f \in C[0,1]$. But these operators are not suitable for discontinuous functions. Later on, Kantorovich[10] generalized the operators(1) to approximate the measurable functions (see G.G. Lorenz[12]). For $n \in N$ and $f \in L_p[0,1], 1 \leq p < \infty$, the Kantorovich operators $K_n : L_p([0,1]) \to L_p([0,1])$ defied by

$$K_n(f;x) = (n+1) \sum_{k=0}^{n} P_{n,k}(x) \int_{\frac{k}{n+1}}^{\frac{k+1}{n+1}} f(t)dt, \qquad k = 1,2,3... \tag{2}$$

Kantorovich type operators converges almost everywhere to the function f on [0,1]. A. Wafi and S. Khatoon[17] defined a generalized Baskakov-Kantorovich operators on $[0,\infty)$

$$V_n^a(f;x) = n \sum_{k=0}^{\infty} W_{n,k}^a(x) \int_{\frac{k}{n}}^{\frac{k+1}{n}} f(t)dt, \tag{3}$$

where

$$W_{n,k}^a(x) = e^{-\frac{ax}{1+x}} \frac{p_k(n,a)}{k!} \frac{x^k}{(1+x)^{k+n}} \tag{4}$$

---

*Corresponding Author
Email addresses: `abdulwafi2k2@gmail.com` (Abdul wafi), `nadeemrao1990@gmail.com` (Nadeem Rao)

Preprint submitted to Name of Journal *August 6, 2015*

such that $\sum\limits_{k=0}^{\infty} W_{n,k}^{a}(x) = 1$ and $p_k(n,a) = \sum\limits_{i=0}^{\infty} \binom{n}{i} (n)_i a^{k-i}$, with $(n)_0 = 1, (n)_i = n(n+1)...(n+i-1)$.

The Stancu operators[14] and its Kantorovich version [2] are respectively given by

$$B_n^{\alpha,\beta}(f;x) = (n+\beta+1)\sum_{k=0}^{n} x^k(1-x)^{n-k} f\left(\frac{k+\alpha}{n+\beta}\right) \tag{5}$$

$$T_n^{\alpha,\beta}(f;x) = (n+\beta+1)\sum_{k=0}^{n} x^k(1-x)^{n-k} \int\limits_{\frac{k+\alpha}{n+\beta+1}}^{\frac{k+\alpha+1}{n+\beta+1}} f(t)dt, \tag{6}$$

where α, β are any two two non negative real numbers such that $0 \le \alpha \le \beta$. If $\alpha = \beta = 0$, the operators (5) and (6) reduce to operators (1) and (2) respectively. In the literature there are many studies on these type of operators(see [1], [2], [7], [8], [11], [15]). In this article, we define generalized Baskakov-Kantorovich-Stancu type opeators as follows

$$T_{n,a}^{\alpha,\beta}(f;x) = (n+\beta)\sum_{k=0}^{\infty} W_{n,k}^{a}(x) \int\limits_{\frac{k+\alpha}{n+\beta}}^{\frac{k+\alpha+1}{n+\beta}} f(t)dt, \tag{7}$$

where $W_{n,k}^{a}(x)$ is defined in (4).

For $\alpha = \beta = 0$, we get the operators (3). The aim of this article is to present a Kantorovich form of Stancu variant of generaliged Baskakov operators (see Rao. N., Wafi A.[13]). Further, we study the direct result, local approximation weighted Korovkin type theorem and order of approximation using weighted modulus of continuity.

2. Basic results

Lemma 2.1 Let $a, x \ge 0$ and $n = 1,2,3....$ Then from[13], we have

$$\begin{aligned}
(i)\quad L_{n,a}^{\alpha,\beta}(1;x) &= 1,\\
(ii)\quad L_{n,a}^{\alpha,\beta}(t;x) &= \frac{n}{n+\beta}x + \frac{a}{n+\beta}\frac{x}{1+x} + \frac{\alpha}{n+\beta},\\
(iii)\quad L_{n,a}^{\alpha,\beta}(t^2;x) &= \frac{n^2+n}{(n+\beta)^2}x^2 + \frac{n(1+2\alpha)}{(n+\beta)^2}x + \frac{a^2}{(n+\beta)^2}\frac{x^2}{(1+x)^2} + \frac{2an}{(n+\beta)^2}\frac{x^2}{(1+x)} + \frac{a(1+2\alpha)}{(n+\beta)^2}\frac{x}{1+x} + \frac{\alpha^2}{(n+\beta)^2}\\
(iv)\quad L_{n,a}^{\alpha,\beta}(t^3;x) &= \frac{n^3+3n^2+2n}{(n+\beta)^3}x^3 + \frac{n^2(3+3\alpha)+n(3+3\alpha+3a)}{(n+\beta)^3}x^2 + \frac{n(1+3\alpha+3\alpha^2)}{(n+\beta)^3}x + \frac{3an^2}{(n+\beta)^3}\frac{x^3}{(1+x)}\\
&\quad + \frac{n}{(n+\beta)^3}\left(\frac{3a^2x^3}{(1+x)^2} + \frac{3ax^2}{1+x} + \frac{6a\alpha x^2}{1+x}\right) + \frac{1}{(n+\beta)^3}\left(\frac{ax}{1+x} + \frac{3a^2x^2}{(1+x)^2} + \frac{a^3x^3}{(1+x)^3}\right.\\
&\quad \left. + \frac{3\alpha a^2x^2}{(1+x)^2} + \frac{3\alpha^2 ax}{1+x} + \alpha^3\right),\\
(v)\quad L_{n,a}^{\alpha,\beta}(t^4;x) &= \frac{n^4+6n^3+11n^2+6n}{(n+\beta)^4}x^4 + \frac{(6+4\alpha)n^3+(18+12\alpha)n^2+(9+8\alpha)n}{(n+\beta)^4}x^3 + \left(\frac{(7+12\alpha+6\alpha^2)n^2}{(n+\beta)^4}\right.\\
&\quad \left. + \frac{(7+12\alpha+12\alpha a+6\alpha^2)n}{(n+\beta)^4}\right)x^2 + \frac{(1+4\alpha+6\alpha^2+4\alpha^3)n}{(n+\beta)^4}x + \frac{4an^3+12an^2+8an}{(n+\beta)^4}\frac{x^4}{1+x}\\
&\quad + \frac{6a^2n^2+6a^2n}{(n+\beta)^4}\frac{x^4}{(1+x)^2} + \frac{4a^3n}{(n+\beta)^4}\frac{x^4}{(1+x)^3} + \frac{a^4}{(b+\beta)^4}\frac{x^4}{(1+x)^4} + \frac{18an^2+18an}{(n+\beta)^4}\frac{x^3}{(1+x)}\\
&\quad + \frac{(18a^2+12a^2\alpha)n}{(n+\beta)^4}\frac{x^3}{(1+x)^2} + \frac{6a^3+4\alpha a^3}{(n+\beta)^4}\frac{x^3}{(1+x)^3} + \frac{(12a\alpha^2+12a\alpha+14a)n}{(n+\beta)^4}\frac{x^2}{(1+x)}\\
&\quad + \frac{7a^2+12a^2\alpha+6a^2\alpha^2}{(n+\beta)^4}\frac{x^2}{(1+x)^2} + \frac{a+4\alpha a+6\alpha^2 a+4\alpha^3 a}{(n+\beta)^4}\frac{x}{1+x} + \frac{\alpha^4}{(n+\beta)^4}.
\end{aligned}$$

Lemma 2.2 Let $a, x \geq 0$. Then

$$\begin{aligned}
(i) \quad T_{n,a}^{\alpha,\beta}(1;x) &= 1, \\
(ii) \quad T_{n,a}^{\alpha,\beta}(t;x) &= \frac{n}{n+\beta}x + \frac{a}{n+\beta}\frac{x}{1+x} + \frac{2\alpha+1}{2(n+\beta)}, \\
(iii) \quad T_{n,a}^{\alpha,\beta}(t^2;x) &= \frac{n^2+n}{(n+\beta)^2}x^2 + \frac{n(2+2\alpha)}{(n+\beta)^2}x + \frac{a^2}{(n+\beta)^2}\frac{x^2}{(1+x)^2} + \frac{2an}{(n+\beta)^2}\frac{x^2}{(1+x)} + \frac{a(2+2\alpha)}{(n+\beta)^2}\frac{x}{1+x} + \frac{3\alpha^2+1}{3(n+\beta)^2} \\
(iv) \quad T_{n,a}^{\alpha,\beta}(t^3;x) &= \frac{n^3+3n^2+2n}{(n+\beta)^3}x^3 + \frac{n^2(\frac{9}{2}+3\alpha)+n(\frac{9}{2}+3\alpha+3a)}{(n+\beta)^3}x^2 + \frac{n(\frac{7}{2}+6\alpha+3\alpha^2)}{(n+\beta)^3}x + \frac{3an^2}{(n+\beta)^3}\frac{x^3}{(1+x)} \\
&\quad + \frac{n}{(n+\beta)^3}\left(\frac{3a^2x^3}{(1+x)^2} + \frac{6ax^2}{1+x} + \frac{6a\alpha x^2}{1+x}\right) + \frac{1}{(n+\beta)^3}\left(a\left(\frac{7}{2}+3\alpha+3\alpha a^2\right)\frac{x}{1+x} + \left(\frac{9}{2}a^2+3\alpha a^2\right)\frac{x^2}{(1+x)^2}\right. \\
&\quad \left. + \frac{a^3x^3}{(1+x)^3}\right) + \frac{\alpha^3+\frac{3}{2}\alpha^2+\alpha}{(n+\beta)^3}, \\
(v) \quad T_{n,a}^{\alpha,\beta}(t^4;x) &= \frac{n^4+6n^3+11n^2+6n}{(n+\beta)^4}x^4 + \frac{(8+4\alpha)n^3+(24+12\alpha)n^2+(13+8\alpha)n}{(n+\beta)^4}x^3 + \left(\frac{(15+18\alpha+6\alpha^2)n^2}{(n+\beta)^4}\right. \\
&\quad \left. + \frac{(15+18\alpha++6a+12\alpha a+6\alpha^2)n}{(n+\beta)^4}\right)x^2 + \frac{(6+14\alpha+11\alpha^2+4\alpha^3)n}{(n+\beta)^4}x + \frac{4an^3+12an^2+8an}{(n+\beta)^4}\frac{x^4}{1+x} \\
&\quad + \frac{6a^2n^2+6a^2n}{(n+\beta)^4}\frac{x^4}{(+x)^2} + \frac{4a^3n}{(n+\beta)^4}\frac{x^4}{(1+x)^3} + \frac{a^4}{(b+\beta)^4}\frac{x^4}{(1+x)^4} + \frac{24an^2+18an}{(n+\beta)^4}\frac{x^3}{(1+x)} \\
&\quad + \frac{(24a^2+12a^2\alpha)n}{(n+\beta)^4}\frac{x^3}{(1+x)^2} + \frac{8a^3+4\alpha a^3}{(n+\beta)^4}\frac{x^3}{(1+x)^3} + \frac{(12a\alpha^2+10a+24a\alpha+14a)n}{(n+\beta)^4}\frac{x^2}{(1+x)} \\
&\quad + \frac{15a^2+18a^2\alpha+6a^2\alpha^2}{(n+\beta)^4}\frac{x^2}{(1+x)^2} + \frac{6a+8\alpha a+12\alpha^2 a+4\alpha^3 a}{(n+\beta)^4}\frac{x}{1+x} + \frac{\alpha^4+2\alpha^3+2\alpha^2+\alpha}{(n+\beta)^4}.
\end{aligned}$$

Proof Identity (i) is obvious. For identity (ii), we use the linearity property

$$T_{n,a}^{\alpha,\beta}(t;x) = (n+\beta)\sum_{k=0}^{\infty} W_{n,k}^{a}(x) \int_{\frac{k+\alpha}{n+\beta}}^{\frac{k+\alpha+1}{n+\beta}} t\,dt,$$

$$T_{n,a}^{\alpha,\beta}(f;x) = (n+\beta)\sum_{k=0}^{\infty} W_{n,k}^{a}(x) \left[\frac{t^2}{2}\right]_{\frac{k+\alpha}{n+\beta}}^{\frac{k+\alpha+1}{n+\beta}},$$

$$T_{n,a}^{\alpha,\beta}(t;x) = \sum_{k=0}^{\infty} W_{n,k}^{a}(x) \left[\frac{k+\alpha}{n+\beta} + \frac{1}{2(n+\beta)}\right].$$

From Lemma 2.1, we have

$$T_{n,a}^{\alpha,\beta}(t;x) = L_{n,a}^{\alpha,\beta}(t;x) + \frac{1}{2(n+\beta)} L_{n,a}^{\alpha,\beta}(1;x)$$

$$T_{n,a}^{\alpha,\beta}(t;x) = \frac{n}{n+\beta}x + \frac{a}{n+\beta}\frac{x}{1+x} + \frac{\alpha}{n+\beta} + \frac{1}{2(n+\beta)}$$

which prove the identity (ii). Similarly, we can prove identities (iii), (iv) and (v).

Lemma 2.3 Let $\psi_x^i(t)=(t-x)^i, i=1,2,3,....$ Then, we have

$$\begin{aligned}
T_{n,a}^{\alpha,\beta}(\psi_x^0(t);x) &= 1,\\
T_{n,a}^{\alpha,\beta}(\psi_x^1(t);x) &= \left(\frac{n}{n+\beta}-1\right)x+\frac{a}{n+\beta}\frac{x}{1+x}+\frac{2\alpha+1}{2(n+\beta)},\\
T_{n,a}^{\alpha,\beta}(\psi_x^2(t);x) &= \frac{n+\beta^2}{(n+\beta)^2}x^2+\frac{n-2(\alpha+1)\beta}{(n+\beta)^2}x+\frac{a^2}{(n+\beta)^2}\frac{x^2}{(1+x)^2}-\frac{2a\beta}{(n+\beta)^2}\frac{x^2}{(1+x)}+\frac{a(2+2\alpha)}{(n+\beta)^2}\frac{x}{1+x}+\frac{3\alpha^2+1}{3(n+\beta)^2},\\
T_{n,a}^{\alpha,\beta}(\psi_x^4(t);x) &= \frac{(3-12\beta)n^2+(6+4\beta+2\beta^2+4\beta^3)n+\beta^4}{(n+\beta)^4}x^4+\left(\frac{(6-12a-12\beta\alpha)n^2+(13+8\alpha-18\beta+12a\beta+9\beta^2)n}{(n+\beta)^4}\right.\\
&\quad\left.+\frac{-4\beta^3\alpha-2\beta^2}{(n+\beta)^4}\right)x^3+\frac{(11-18\alpha+18\alpha^2)n^2+(15+18\alpha+6a+12\alpha a-24\alpha\beta)n+6\alpha^2\beta^2+2\beta^2}{(n+\beta)^4}x^2\\
&\quad+\frac{(6+10\alpha+5\alpha^2)n-4\beta(\alpha^3+\frac{3}{2}\alpha^2+\alpha)}{(n+\beta)^4}x+\frac{12an^2+8an-4a\beta^3}{(n+\beta)^4}\frac{x^4}{(1+x)}\\
&\quad+\frac{6a^2n+6a^2\beta^2}{(n+\beta)^4}\frac{x^4}{(1+x)^2}-\frac{4a^3\beta}{(n+\beta)^4}\frac{x^4}{(1+x)^3}+\frac{a^4}{(n+\beta)^4}\frac{x^4}{(1+x)^4}+\frac{12an^2+18an+6a(1+2\alpha)\beta^2}{(n+\beta)^4}\frac{x^3}{1+x}\\
&\quad+\frac{6a^2n-(12a^2+12\alpha a^2)\beta}{(n+\beta)^4}\frac{x^3}{(1+x)^2}+\frac{6a^3+4\alpha a^3}{(n+\beta)^4}\frac{x^3}{(1+x)^3}+\frac{(12a\alpha+8a-6a\alpha^2)n-(6a+18\alpha^2a)\beta}{(n+\beta)^4}\frac{x^2}{1+x}\\
&\quad+\frac{7a^2+12a^2\alpha+6a^2\alpha^2}{(n+\beta)^4}\frac{x^2}{(1+x)^2}+\frac{(a)+4\alpha a+6\alpha^2a+4\alpha^3a}{(n+\beta)^4}\frac{x}{1+x}+\frac{\alpha^4}{(n+\beta)^4}.
\end{aligned}$$

Proof By using the linearity property and Lemma 2.2, we can prove the Lemma2.3 .

Lemma 2.4 For the operators $T_{n,a}^{\alpha,\beta}$, we have

$$T_{n,a}^{\alpha,\beta}((t-x)^4;x) \quad\leq\quad M_a^{\alpha,\beta}\frac{(x^4+x^3+x^2+x+1)}{(n+\beta)^2}$$

where $M_a^{\alpha,\beta}$ is a positive constant for the fixed value of non-negative numbers $a,\alpha,$ and β.

Proof Since $\frac{x^i}{(1+x)^j}\leq x^i$ for all $x\geq 0$ and $j\leq i(j,i=1,2,3,4)$, we have

$$\begin{aligned}
T_{n,a}^{\alpha,\beta}((t-x)^4;x) &\leq \frac{1}{(n+\beta)^2}\left(\frac{(3-12\beta+12a)n^2+(6+4\beta+2\beta^2+4\beta^3+8a+6a^2)n+\beta^4-4a\beta^3+6a^2\beta^2-4a^3\beta+a^4}{(n+\beta)^2}x^4\right.\\
&\quad+\left(\frac{(6-12a-12\beta\alpha+12a)n^2+(13+8\alpha-18\beta+12a\beta+9\beta^2+18a+6a^2)n-4\beta^3\alpha-2\beta^2}{(n+\beta)^2}\right.\\
&\quad\left.\frac{(6a(1+2\alpha)\beta^2-12a^2+12\alpha a^2)\beta+6a^3+4\alpha a^3}{(n+\beta)^4}\right)x^3+\left(\frac{(11-18\alpha+18\alpha^2)n^2}{(n+\beta)^2}\right.\\
&\quad\left.+\frac{(15+18\alpha+14a+24\alpha a-24\alpha\beta-6a\alpha^2)n+12\alpha^2\beta^2+2\beta^2-(6a+18\alpha^2a)\beta+7a^2+12a^2\alpha}{(n+\beta)^2}\right)x^2\\
&\quad+\frac{(6+10\alpha+5\alpha^2)n-4\beta(\alpha^3+\frac{3}{2}\alpha^2+\alpha)+a+4\alpha a+6\alpha^2a+6\alpha^3a}{(n+\beta)^2}x,
\end{aligned}$$

$$T_{n,a}^{\alpha,\beta}((t-x)^4;x)\leq\frac{A_4x^4+A_3x^3+A_2x^2+A_1x}{(n+\beta)^2} \tag{8}$$

where

$$\begin{aligned}
A_4 &= \frac{(3-12\beta+12a)n^2+(6+4\beta+2\beta^2+4\beta^3+8a+6a^2)n+\beta^4-4a\beta^3+6a^2\beta^2-4a^3\beta+a^4}{(n+\beta)^2}\\
A_3 &= \frac{(6-12a-12\beta\alpha+12a)n^2+(13+8\alpha-18\beta+12a\beta+9\beta^2+18a+6a^2)n-4\beta^3\alpha-2\beta^2}{(n+\beta)^2}\\
&\quad +\frac{(6a(1+2\alpha)\beta^2-12a^2+12\alpha a^2)\beta+6a^3+4\alpha a^3}{(n+\beta)^4}\\
A_2 &= \frac{(15+18\alpha+14a+24\alpha A-24\alpha\beta-6a\alpha^2)n+12\alpha^2\beta^2+2\beta^2-(6a+18\alpha^2 a)\beta+7a^2+12a^2\alpha}{(n+\beta)^2}\Bigg)\\
A_1 &= \frac{(6+10\alpha+5\alpha^2)n-4\beta(\alpha^3+\frac{3}{2}\alpha^2+\alpha)+a+4\alpha a+6\alpha^2 a+6\alpha^3 a}{(n+\beta)^2}.
\end{aligned}$$

For large value of n, each $A_i (i=1,2,3,4)$ converges to a value depending on a,α,β, Therefore, there exists a constant $M_a^{\alpha,\beta}$ for the fixed values of a,α,β such that $A_i \leq M_a^{\alpha,\beta}$ for $i=1,2,3,4$. Using (8), we arrive at the result.

3. Direct approximation

Let $C_B[0,\infty)$ denote the space of real valued continuous and bounded functions f on $[0,\infty)$ endowed with the norm

$$\|f\| = \sup_{0\leq x<\infty} |f(x)|.$$

Then, for any $\delta > 0$, Peeter's K-functional is defined as

$$K_2(f,\delta) = inf\{\|f-g\|+\delta\|g''\| : g \in C_B^2[0,\infty)\},$$

where $C_B^2[0,\infty) = \{g \in C_B[0,\infty) : g', g'' \in C_B[0,\infty)\}$. By Devore and Lorentz[[4], p.177, Theorem 2.4], there exits an absolute constant $C > 0$ such that

$$K_2(f;\delta) \leq C\omega_2(f;\sqrt{\delta}),$$

where $\omega_2(f;\delta)$ is the second order modulus of continuity is defined as

$$\omega_2(f,\sqrt{\delta}) = \sup_{0<h\leq\sqrt{\delta}} \sup_{x\in[0,\infty)} |f(x+2h)-2f(x+h)+f(x)|.$$

Theorem 3.1 Let $f \in C_B^2[0,\infty)$. Then for all $x \in [0,\infty)$ there exist a constant $K > 0$ such that

$$| T_{n,a}^{\alpha,\beta}(f;x)-f(x) | \leq K\omega_2(f;\sqrt{\gamma_n^{\alpha,\beta}(x)})+\omega\left(f;\left(\frac{\beta}{n+\beta}\right)x+\frac{a}{n+\beta}\frac{x}{1+x}+\frac{2\alpha+1}{2(n+\beta)}\right)$$

where $\gamma_n^{\alpha,\beta}(x) = \left\{\frac{n+2\beta^2}{(n+\beta)^2}x^2+\frac{n-\beta}{(n+\beta)^2}x+\frac{2a^2}{(n+\beta)^2}\frac{x^2}{(1+x)^2}+\frac{a(3+4\alpha)}{(n+\beta)^2}\frac{x}{1+x}+\frac{7\alpha^2+4\alpha+2}{3(n+\beta)^2}\right\}$.

Proof First, we define the auxiliary operators

$$\hat{T}_n^{\alpha,\beta}(f;x) = T_{n,a}^{\alpha,\beta}(f;x)+f(x)-f\left(\frac{n}{n+\beta}x+\frac{a}{n+\beta}\frac{x}{1+x}+\frac{2\alpha+1}{2(n+\beta)}\right) \tag{9}$$

We find that

$$\hat{T}_n^{\alpha,\beta}(1;x) = 1,$$

$$\hat{T}_n^{\alpha,\beta}(\psi_x(t);x) = 0$$

$$|\hat{T}_n^{\alpha,\beta}(f;x)| \leq 3\|f\|. \tag{10}$$

Let $g \in C_B^2[0,\infty)$. By the Taylor's theorem

$$g(t) = g(x) + (t-x)g'(x) + \int\limits_x^t (t-v)g''(v)dv, \tag{11}$$

Now, using (11), the auxiliary operators is given

$$\begin{aligned}
\hat{T}_n^{\alpha,\beta}(g;x) - g(x) &= g'(x)\hat{T}_n^{\alpha,\beta}(t-x;x) + \hat{T}_n^{\alpha,\beta}\Big(\int\limits_x^t (t-v)g''(v)dv;x\Big)\\
&= \hat{T}_n^{\alpha,\beta}\Big(\int\limits_x^t (t-v)g''(v)dv;x\Big)\\
&= T_{n,a}^{\alpha,\beta}\Big(\int\limits_x^t (t-v)g''(v)dv;x\Big) - \int\limits_x^{\frac{n}{n+\beta}x+\frac{a}{n+\beta}\frac{x}{1+x}+\frac{2\alpha+1}{2(n+\beta)}} \Big(\frac{n}{n+\beta}x + \frac{a}{n+\beta}\frac{x}{1+x} + \frac{2\alpha+1}{2(n+\beta)} - v\Big)g''(v)dv.
\end{aligned}$$

Therefore

$$\begin{aligned}
|\hat{T}_n^{\alpha,\beta}(g;x) - g(x)| &\leq \left|T_{n,a}^{\alpha,\beta}\Big(\int\limits_x^t (t-v)g''(v)dv;x\Big)\right|\\
&\quad + \left|\int\limits_x^{\frac{n}{n+\beta}x+\frac{a}{n+\beta}\frac{x}{1+x}+\frac{2\alpha+1}{2(n+\beta)}} \Big(\frac{n}{n+\beta}x + \frac{a}{n+\beta}\frac{x}{1+x} + \frac{2\alpha+1}{2(n+\beta)} - v\Big)g''(v)dv\right|.
\end{aligned} \tag{12}$$

Since

$$\left|\int\limits_x^t (t-v)g''(v)dv\right| \leq (t-x)^2 \parallel g'' \parallel \tag{13}$$

and

$$\left|\int\limits_x^{\frac{n}{n+\beta}x+\frac{a}{n+\beta}\frac{x}{1+x}+\frac{2\alpha+1}{2(n+\beta)}} \Big(\frac{n}{n+\beta}x + \frac{a}{n+\beta}\frac{x}{1+x} + \frac{2\alpha+1}{2(n+\beta)} - v\Big)g''(v)dv\right| \leq \left(\frac{\beta}{n+\beta}x + \frac{a}{n+\beta}\frac{x}{1+x} + \frac{2\alpha+1}{2(n+\beta)}\right)^2 \parallel g'' \parallel. \tag{14}$$

Then from(12),(13)and (14) implies that

$$\begin{aligned}
|\hat{T}_n^{\alpha,\beta}(g;x) - g(x)| &\leq \left\{T_{n,a}^{\alpha,\beta}((t-x)^2;x) + \left(\frac{\beta}{n+\beta}x + \frac{a}{n+\beta}\frac{x}{1+x} + \frac{2\alpha+1}{2(n+\beta)}\right)^2\right\}\|g''\|\\
&= \left\{\frac{n+2\beta^2}{(n+\beta)^2}x^2 + \frac{n-\beta}{(n+\beta)^2}x + \frac{2a^2}{(n+\beta)^2}\frac{x^2}{(1+x)^2} + \frac{a(3+4\alpha)}{(n+\beta)^2}\frac{x}{1+x} + \frac{7\alpha^2+4\alpha+2}{3(n+\beta)^2}\right\}\|g''\|\\
&= \gamma_n^{\alpha,\beta}(x)\|g''\|
\end{aligned} \tag{15}$$

Next, we have

$$|T_{n,a}^{\alpha,\beta}(f;x) - f(x)| \leq |\hat{T}_n^{\alpha,\beta}(f-g;x)| + |(f-g)(x)| + |\hat{T}_n^{\alpha,\beta}(g;x) - g(x)| + \big|f(\frac{n}{n+\beta}x + \frac{a}{n+\beta}\frac{x}{1+x} + \frac{2\alpha+1}{2(n+\beta)}) - f(x)\big|$$

Using(15), we have

$$\begin{aligned}
|T_{n,a}^{\alpha,\beta}(f;x) - f(x)| &\leq 4\|f-g\| + \hat{T}_n^{\alpha,\beta}(g;x) - g(x)| + \big|f(\frac{n}{n+\beta}x + \frac{a}{n+\beta}\frac{x}{1+x} + \frac{2\alpha+1}{2(n+\beta)}) - f(x)\big|\\
&\leq 4\|f-g\| + \gamma_n^{\alpha,\beta}(x)\|g''\| + \omega\Big(f;\frac{n}{n+\beta}x + \frac{a}{n+\beta}\frac{x}{1+x} + \frac{2\alpha+1}{2(n+\beta)}\Big)
\end{aligned}$$

By the definition of Peetre's K-functional

$$|T_{n,a}^{\alpha,\beta}(f;x)-f(x)| \le C\omega_2\big(f;\sqrt{\gamma_n^{\alpha,\beta}(x)}\big)+\omega\Big(f;\frac{n}{n+\beta}x+\frac{a}{n+\beta}\frac{x}{1+x}+\frac{2\alpha+1}{2(n+\beta)}\Big).$$

Now, we use the Lipschitz type space

$$LIp_M^* = \{f\in C[0,\infty): |f(t)-f(x)| \le M\frac{|t-x|^\alpha}{(t+x)^{\frac{\alpha}{2}}} : x,t\in(0,\infty)\}$$

where M is a constant and $0<\alpha\le 1$ to prove the following theorem:

Theorem 3.2 Let $f\in Lip_M^*(\alpha)$ and $x\in[0,\infty)$. Then, we have

$$|T_{n,a}^{\alpha,\beta}(f;x)-f(x)| \le M\left[\frac{\Lambda_n^{\alpha,\beta}(x)}{x}\right],$$

where $\Lambda_n^{\alpha,\beta}(x)=T_n^{\alpha,\beta}((t-x)^2;x)$.

Proof Let $\alpha=1$ and $x\in(0,\infty)$. Then, for $f\in Lip_M^*(1)$, we have

$$\begin{aligned}
|T_{n,a}^{\alpha,\beta}(f;x)-f(x)| &\le (n+\beta)\sum_{k=0}^{\infty}W_{n,k}^a(x)\int_{\frac{k+\alpha}{n+\beta}}^{\frac{k+\alpha+1}{n+\beta}}|f(t)-f(x)|dt \\
&\le M(n+\beta)\sum_{k=0}^{\infty}W_{n,k}^a(x)\int_{\frac{k+\alpha}{n+\beta}}^{\frac{k+\alpha+1}{n+\beta}}\frac{|t-x|}{\sqrt{t+x}}dt. \\
&\le \frac{M}{\sqrt{x}}(n+\beta)\sum_{k=0}^{\infty}W_{n,k}^a(x)\int_{\frac{k+\alpha}{n+\beta}}^{\frac{k+\alpha+1}{n+\beta}}|t-x|dt \\
&\le \frac{M}{\sqrt{x}}T_n^{\alpha,\beta}(|t-x|;x) \\
&\le M\frac{\sqrt{T_n^{\alpha,\beta}((t-x)^2;x)}}{\sqrt{x}} \\
&= M\left(\frac{\Lambda_n^{\alpha,\beta}(x)}{x}\right)^{\frac{1}{2}}
\end{aligned}$$

Thus, the assertion hold for $\alpha=1$. Now, we will prove for $\alpha\in(0,1)$. From the Holder inequality with $p=\frac{1}{\alpha}, q=\frac{1}{1-\alpha}$, we have

$$\begin{aligned}
|T_{n,a}^{\alpha,\beta}(f;x)-f(x)| &= \left(\sum_{k=0}^{\infty}W_{n,k}^a(x)\left((n+\beta)\int_{\frac{k+\alpha}{n+\beta}}^{\frac{k+\alpha+1}{n+\beta}}|f(t)-f(x)|dt\right)^{\frac{1}{\alpha}}\right)^{\alpha}\left(W_{n,k}^a(x)\right)^{1-\alpha} \\
&\le \left(\sum_{k=0}^{\infty}W_{n,k}^a(x)\left((n+\beta)\int_{\frac{k+\alpha}{n+\beta}}^{\frac{k+\alpha+1}{n+\beta}}|f(t)-f(x)|dt\right)^{\frac{1}{\alpha}}\right)^{\alpha}
\end{aligned}$$

Since $f\in Lip_M^*$, we obtain

$$\begin{aligned}
|T_{n,a}^{\alpha,\beta}(f;x)-f(x)| &\le M\left((n+\beta)\sum_{k=0}^{\infty}W_{n,k}^a(x)(\int_{\frac{k+\alpha}{n+\beta}}^{\frac{k+\alpha+1}{n+\beta}}\frac{|t-x|}{\sqrt{t+x}}dt\right)^{\alpha} \\
&\le \frac{M}{x^{\frac{\alpha}{2}}}\left((n+\beta)\sum_{k=0}^{\infty}W_{n,k}^a(x)\int_{\frac{k+\alpha}{n+\beta}}^{\frac{k+\alpha+1}{n+\beta}}|t-x|dt\right)^{\alpha} \\
&= \frac{M}{x^{\frac{\alpha}{2}}}\left(T_{n,a}^{\alpha,\beta}(|t-x|;x)\right)^{\alpha} \\
&\le M\left(\frac{\Lambda_n^{\alpha,\beta}(x)}{x}\right)^{\frac{r}{2}}.
\end{aligned}$$

4. Approximation properties of $T_{n,a}^{\alpha,\beta}$ in weighted space

In this section, we investigate the weighted approximation properties of the operators $T_{n,a}^{\alpha,\beta}$ by the weighted Korovkin type theorem given by Gadzhiev ([5], [6]) and using weighted modulus of continuity defined by Ispir[9].

Let $B_\rho[0,\infty) = \{f(x) : |f(x)| \leq M_f \rho(x), \rho(x)$ is weight function, M_f is a constant depending on f and $x \in [0,\infty)\}$, $C_\rho[0,\infty)$ is the space of continuous function in $B_\rho[0,\infty)$ with the norm $\|f(x)\|_\rho = \sup\limits_{x\in[0,\infty)} \frac{|f(x)|}{\rho(x)}$ and $C_\rho^k = \{f \in C_\rho : \lim\limits_{|x|\to\infty} \frac{f(x)}{\rho(x)} = k$, where k is a constant depending on $f\}$.

Theorem 4.1 Let $T_{n,a}^{\alpha,\beta}$ be the sequence of linear positive operators defined by (7). Then for $f \in C_\rho^k$,

$$\lim_{n\to\infty} \|T_{n,a}^{\alpha,\beta}(f;x) - f(x)\|_\rho = 0.$$

Proof To prove the theorem, it is sufficient to show that

$$\lim_{n\to\infty} \|T_{n,a}^{\alpha,\beta}(t^i;x) - x^i\|_\rho = 0, \qquad for \quad i = 0,1,2.$$

It is obvious that $\lim\limits_{n\to\infty} \|T_{n,a}^{\alpha,\beta}(1;x) - 1\|_\rho = 0$. Now,from the Lemma 2.2, we have

$$\begin{aligned}\|T_{n,a}^{\alpha,\beta}(t;x) - x\| &= \sup_{x\in[0,\infty)} \frac{\left|\left(\frac{n}{n+\beta} - 1\right)x + \frac{a}{n+\beta}\frac{x}{1+x} + \frac{2\alpha+1}{2(n+\beta)}\right|}{1+x^2} \\ &\leq \left(\frac{\beta}{n+\beta}\right) \sup_{x\in[0,\infty)} \frac{x}{1+x^2} + \frac{a}{n+\beta} \sup_{x\in[0,\infty)} \frac{x}{(1+x)(1+x)^2} + \frac{2\alpha+1}{2(n+\beta)} \sup_{x\in[0,\infty)} \frac{1}{(1+x)^2}.\end{aligned}$$

$$\| \hat{T}_n^{\alpha,\beta}(t;x) - x \|_\rho \to 0 \qquad as \;\; n \to \infty.$$

Also, we can write

$$\begin{aligned}\|T_{n,a}^{\alpha,\beta}(t^2;x) - x^2\| &= \sup_{x\in[0,\infty)} \frac{\left|\frac{n^2+n}{(n+\beta)^2}x^2 + \frac{n(2+2\alpha)}{(n+\beta)^2}x + \frac{a^2}{(n+\beta)^2}\frac{x^2}{(1+x)^2} + \frac{2an}{(n+\beta)^2}\frac{x^2}{(1+x)} + \frac{a(2+2\alpha)}{(n+\beta)^2}\frac{x}{1+x} + \frac{3\alpha^2+1}{3(n+\beta)^2} - x^2\right|}{1+x^2} \\ &\leq \frac{n(1-2\beta)+\beta^2}{(n+\beta)^2} \sup_{x\in[0,\infty)} \frac{x^2}{1+x^2} + \frac{n(2+2\alpha)}{(n+\beta)^2} \sup_{x\in[0,\infty)} \frac{x}{1+x^2} + \frac{a^2}{(n+\beta)^2} \sup_{x\in[0,\infty)} \frac{x^2}{(1+x)^2(1+x^2)} \\ &\quad + \frac{2an}{(n+\beta)^2} \sup_{x\in[0,\infty)} \frac{x^2}{(1+x)(1+x^2)} + \frac{a(2+2\alpha)}{(n+\beta)^2} \sup_{x\in[0,\infty)} \frac{x}{(1+x)(1+x^2)} + \frac{3\alpha^2+1}{3(n+\beta)^2} \sup_{x\in[0,\infty)} \frac{1}{(1+x^2)}.\end{aligned}$$

Which implies that $\|T_{n,a}^{\alpha,\beta}(t^2;x) - x^2\|_\rho \to 0$ as $n \to \infty$. Hence, proof is completed.

Now we prove the next theorem using weighted modulus of continuity defined by Ispir[9] as:

$$\Omega_n(f;\delta) = \sup_{|h|\leq\delta, x\in[0,\infty)} \frac{|f(x+h) - f(x)|}{(1+h^2)(1+x^2)},$$

for each $f \in C_\rho^k[0,\infty)$.
Some properties of $\Omega_n(f;\delta)$ are as follows:

Lemma 4.2 Let $f \in C_\rho^K$,

(i) $\Omega_n(f;\delta)$ is a monotonically increasing function of δ, $\delta \geq 0$.

(ii) For every $f \in C_\rho^k$, $\lim\limits_{\delta\to 0} \Omega_n(f;\delta) = 0$.

(iii) For each positive value of λ

$$\Omega_n(f;\lambda\delta) \leq 2(1+\lambda)(1+\delta^2)\Omega_n(f;\delta).$$

Using this property of modulus of continuity $\Omega_n(f;\delta)$ and its definition, we have

$$|f(t)-f(x)| \leq (1+x^2)(1+(t-x)^2)\Omega_n(f;|t-x|)$$

and also

$$\begin{aligned} |f(t)-f(x)| &\leq 2 \quad \Big(\frac{|t-x|}{\delta_n}+1\Big)\Omega_n(f;\delta_n)(1+x^2)(1+(t-x)^2). && (16)\\ \Big(\frac{|t-x|}{\delta_n}+1\Big)(1+(t-x)^2) &\leq 2 \quad (1+\delta_n^2)\Big(1+\frac{(t-x)^4}{\delta_n^4}\Big) && (17) \end{aligned}$$

Theorem 4.3 Let $f \in C^k_{\rho,[0,\infty)}$. Then the inequality

$$\sup_{x\in[0,\infty)} \frac{|T^{\alpha,\beta}_{n,a}(f;x)-f(x)|}{(1+x^2)^3} \leq M_a^{\alpha,\beta}\Omega_n(f;(n+\beta)^{-\frac{1}{2}})$$

holds, where $M_a^{\alpha,\beta}$ is a constant independent of n.

Proof From (16) and (17) we have

$$\begin{aligned} |T^{\alpha,\beta}_{n,a}(f;x)-f(x)| &\leq 4(1+x^2)(1+\delta_n^2)\Omega_n(f;\delta_n)T^{\alpha,\beta}_{n,a}\Big(1+\frac{(t-x)^4}{\delta_n^4}\Big),\\ &= 4(1+x^2)(1+\delta_n^2)\Omega_n(f;\delta_n)\Big(1+T^{\alpha,\beta}_{n,a}\Big(\frac{(t-x)^4}{\delta_n^4};x\Big)\Big)\\ &\leq (1+x^2)\Omega_n(f;\delta_n)\Big(1+\frac{1}{\delta^4}T^{\alpha,\beta}_{n,a}((t-x)^4;x). \end{aligned}$$

Using Lemma 2.4, we obtain

$$\begin{aligned} |T^{\alpha,\beta}_{n,a}(f;x)-f(x)| &\leq M_a^{\alpha,\beta}(1+x^2)\Omega_n(f;\delta_n)\Big(1+\frac{\Big(\frac{1}{(n+\beta^2)}\Big)(x^4+x^3+x^2+x+1)}{\delta_n^4}\Big)\\ &\leq M_a^{\alpha,\beta}(1+x^2)\Omega_n(f;(n+\beta)^{-\frac{1}{2}})(x^4+x^3+x^2+x+1)\\ &\leq M_a^{\alpha,\beta}(1+x^2)^3\Omega_n(f;(n+\beta)^{-\frac{1}{2}}). \end{aligned}$$

Therefore, we obtain

$$\sup_{x\in[0,\infty)} \frac{|T^{\alpha,\beta}_{n,a}(f;x)-f(x)|}{(1+x^2)^3} \leq M_a^{\alpha,\beta}\Omega_n(f;(n+\beta)^{-\frac{1}{2}}).$$

5. References

References

[1] Aral A., Acar T., Weightet Approximation by new Bernstein -Chlodowsky-Gadjiev operators, Filomat, 2013, **27 (2)** 371-380.

[2] Basrbosu D., Kantorovich-Stancu type operators, J. Inequal. Pure Appl. Math., 2004, **5 (3)**, Article 53, 6 pp.

[3] Bernstein S.N., Dmonstration du thorms de Weierstrass fond sur le calcul des probabilits, Commun. Soc. Math. Kharkow, 1912-13, **13 (2)**, 12.

[4] DeVore R.A., Lorentz G.G., Constructive Approximation, Grudlehren der Mathematischen Wissenschaften [Fundamental principales of Mathematical Sciences], (Springer-Verlag, Berlin, 1993).

[5] Gadjiev A.D., The convergence problem for a sequence of positive linear operators on unbounded sets and theorems analogous to that P.P. Korovkin, Soviet Math. Dokl., 1974, **15 (5)**, 1433-1436.

[6] Gadjiev A.D., Theorems of the type of P.P. Korovkin's theorems, Math. Zametki, 1976, **20 (5)**, 781-786.(in Russian), Math. Notes, 1976, **20 (5-6)**, 995-998(Engl. Trans.).
[7] Gupta V., Agarwal R.P., Convergence Estimates in Approximtion Theory, Springer, Cham,2014.
[8] Gupta V.,Verma D.K., Agrawal P.N., Simultaneous approximation be crtain Baskakov-Stancu Operators,J. Egyptian Math. Soc., 2012, **20 (3)**, 183-187.
[9] Ispir N., On modified Baskakov operators on weighted spaces, Turkish J. Math., 2001, **25 (3)**, 355-365.
[10] Kantorovich L.V., Sur certain developpments suivant les polynomes de la forme de S. Bernstein, I,II, C.R. Acad. URSS, 1930, 563-568, 595-600.
[11] Lopez-Moreno A.-J., Weighted simultaneous approximaton with Baskakov type operators, Acta Math. Hungar., 2004, **104 (1-2)**, 143-151.
[12] Lorentz G.G., Mathematical Expositions, No. 8, Bernstein polynomials, University of Toronto Press, Toronto 1953.
[13] Rao N., Wafi A.; Stancu-varient of generalised Baskakov operators, Filomat, 2015, (Accepted).
[14] Stancu D.D., On a generalization of the Bernstein polynomials(Romanian), Stud. Univ. Babes-Bolyai Math., 1969, **14 (2)**, 31-45.
[15] Sucu S., Ibikli E., Approximation by means of Kantorovich-Stancu type operators, Numer. Funct. Anal. Optim., 2013, **34 (5)**, 557-575.
[16] Wafi A., Khatoon S., The Voronovskaya theorem for generalized Baskakov-Kantorovich operalors in polynomial weight spaces, Matematicki Vesnik, 2005, **57 (3-4)**, 87-94.
[17] Wafi A., Khatoon S., Uniform approximation with generalized Baskakov-Kantorovich operators, Alig. Bull. Math., 2003, **22 (2)**, 119-133.